

DIGESTIVE AND LIVER DISEASE SPECIALISTS

MoviPrep PREPARATION

Purchase MoviPrep from your pharmacy. You will need a prescription.
Begin a clear liquid diet on the morning of _____. (See next page for clear liquid diet)

Beginning at: _____

STEP 1: Empty 1 Pouch A and 1 Pouch B into the disposable container.
Add lukewarm drinking water to the top line of the container. Mix to dissolve.

You may chill the reconstituted solution. The solution should be used within 24 hours.

STEP 2: The Moviprep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is complete.

Beginning at: _____ **Repeat steps 1 and 2.**

Stop Coumadin / Warfarin 3 days prior to your procedure. Notify your cardiologist and ask if he wants to replace these medications with Lovenox during this time.

Stop Plavix 7 days prior to your procedure. Notify your cardiologist for approval.

Stop Aggrenox and Ticlid 5 days prior to your procedure. Notify your cardiologist for approval.

Stop Aspirin and Anti Inflammatory medications (ex. Naprosyn, Motrin) 5 days prior to your procedure.

Stop Iron 7 days prior to your procedure.

Celebrex and Tylenol are allowed.

Adjust your diabetic medications per office protocol.

You may take your heart and blood pressure medicine the morning of the procedure.

Check with your physician if there is any question about medications prior to your procedure.

Nothing by mouth 3 hours prior to your exam unless you need to take medications as directed by your physician. These may be taken with a small sip of water.

DUE TO THE SEDATION, YOU MUST HAVE SOMEONE DRIVE YOU HOME

